



RESEARCH ARTICLE

Exploring Pesticide Application Practices for Managing Prominent Agricultural Pests in and around region of Tenkasi district, Tamil Nadu, India.

Chelliah Ramalakshmi^{1*}, Kandsamy Kalaivani², & Sulaihal Barveen²

¹DMI - St.Eugene University, Chibombo, Zambia.

²Department of Zoology, Sri Parasakthi College for Women, Tenkasi, Tamil Nadu, India.

ARTICLE HISTROY

Received 17 November 2023

Revised 19 November 2023

Accepted 23 November 2023

Keywords

Pesticide

Dimethoate

Ecotoxicology

Integrated pest management

ABSTRACT

This study, conducted between November and March in the Tenkasi district, examined pesticide usage in relation to crop cultivation in Sengottai, Sundarapandiapuram, Sambavar Vadakarai, Panpoli, Kadayam, and Surandai. Sunflower, coconut, raw banana, paddy, and potato were the prominently cultivated crops. Pesticides like neem oil, FAME, carbaryl, coragen, malathion, and dimethoate were employed based on specific crops, showcasing diverse pest management strategies. Analysis revealed Dimethoate as the most extensively used pesticide, underscoring its significance in pest management during the study period. The study highlighted the broader ecological impact of pesticides, emphasizing risks to wildlife and plants. Ecotoxicology, focusing on pesticide impacts, was introduced. The study noted the far-reaching effects of pesticides, considering runoff, wind drift, and the persistence of certain pesticides like DDT and chlordane. Plants' absorption of pesticides through roots and leaves indicated their susceptibility. The cumulative impact on individual plants and animals underscored the need to address potential ecological consequences. In the study urged a careful evaluation of pesticide usage, considering environmental impacts. Adhering to label instructions and exploring alternatives with lower toxicity was advised. Integrated pest management (IPM) and proactive measures to prevent contamination in crucial locations were recommended. The study proposed organic farming as a sustainable alternative, contributing to a healthy environment and human well-being.

✉ Ramalakshmi
ramalakshmispc@gmail.com

©2023 The Author(s). Published by Panainool Ltd.

Introduction

Agriculture is the science and practice of cultivating the soil to grow crops and rear animals for food, wool, and other essential products. One of the primary challenges in agriculture is the



presence of pests that can attack crops, leading to reduced yields (Dhanaraju et al., 2022; Tudi et al., 2021). Pests are defined as any animals or plants that have harmful effects on humans, their food supply, or living conditions. In simpler terms, pests can be described as destructive insects or other animals that damage crops, food, livestock, and more. Examples of pests include aphids, fruit flies, bollworms, whiteflies, leafhoppers, cutworms, beetles, and various other creatures (Ratnadass et al., 2012; Skendžić et al., 2021).

To manage and control these pests, pesticides are employed. Pesticides are substances designed for the purpose of preventing, destroying, or controlling pests and unwanted plant species that can harm or interfere with food production, processing, storage, transportation, or marketing (Pathak et al., 2022; Tudi et al., 2021). The term "pesticide" encompasses various categories, such as herbicides (used to control weeds), nematicides, molluscicides, piscicides, avicides, rodenticides, bactericides, insect repellents, animal repellents, antimicrobials, and fungicides. Among these, herbicides stand out as the most widely used, accounting for 80% of all pesticides (Park et al., 2021).

Pesticides are crucial in agriculture because they protect crops from pests, diseases, and weeds, which, if left unmanaged, could result in the loss of more than half of the world's crops. These chemicals play a vital role in enabling farmers to produce more food with less land, as they enhance productivity per hectare. In the world of pesticides, many substances can be grouped into chemical families. Notable insecticides include

organochlorides, organophosphates, and carbamates (Cooper & Dobson, 2007).

Pesticides are essential tools in modern agriculture and public health, designed to combat a wide range of organisms that can harm crops, livestock, or human well-being. These chemical formulations are created to target and manage pests, which can encompass insects, weeds, fungi, bacteria, and other undesirable organisms. Pesticides are critical in safeguarding agricultural yields, reducing crop loss, and ensuring food security by controlling pests that threaten the quality and quantity of harvests (Nicolopoulou-Stamati et al., 2016).

Pesticides exist in various forms, including insecticides to tackle insect pests, herbicides to suppress weed growth, fungicides to combat fungal diseases, and bactericides to control harmful bacteria. They can be chemically synthesized or derived from natural sources. The use of pesticides has been instrumental in increasing agricultural productivity, helping to feed a growing global population (Abdollahdokht et al., 2022; Ayilara et al., 2023; Hassaan & El Nemr, 2020).

The utilization of pesticides is not without challenges and concerns. Their application may unintentionally harm non-target organisms, lead to environmental contamination, and contribute to the development of pesticide-resistant pests. Striking a balance between the benefits of pest control and minimizing potential risks is an ongoing priority in agriculture and public health. This requires a deep understanding of the different types of pesticides, their mechanisms of action,

and their environmental impacts, with a focus on using them responsibly and effectively.

Materials and Methods

Study area

The research was conducted in the Tenkasi district of Tamil Nadu, focusing on specific regions

including Sengottai, Sundarapandiyapuram, Sambavar Vadakarai, Panpoli, Kadayam, and Surandai (Fig 1). These areas were chosen due to the diverse range of vegetables and crops cultivated throughout the year. The selection of the study area was based on the types of crops grown and the extent of pesticide usage.




Fig 1. Map shows the study sites of Tenkasi District region

The crops cultivated in this region encompass a wide variety, such as potatoes, paddy, drumsticks, raw bananas, coconuts, carrots, onions, tomatoes, brinjal, okra, sunflowers, tapioca, and more. The farmers in these areas are actively engaged in large-scale crop and vegetable cultivation, which has led to a notable and variable increase in pesticide usage over time.

Pesticide use: This section was used to obtain data about the types of pesticides used and sources of pesticides commonly used by the farmers.

Results and discussion

The survey took place from November to March, during which the study site selection was tailored to the crops cultivated in the region. The study area comprised Sengottai, Sundarapandiyapuram, Sambavar Vadakarai, Panpoli, Kadayam, and Surandai in the Tenkasi district, where a diverse array of crops and vegetables were grown. The primary crops included sunflower, coconut, raw banana, paddy, and potato. The usage of pesticides varied depending on the cultivated crops, with

notable pesticides such as neem oil, FAME, carbaryl, coragen, malathion, and dimethoate being employed more extensively based on the specific crops in cultivation. The observation leads to the conclusion that Dimethoate is the predominant pesticide used as shown in Table 1.

Pesticides and other contaminants entering the natural environment have the potential to impact wild plants and animals, a concern addressed by the field of ecotoxicology (Damalas & Eleftherohorinos, 2011; Fleeger, 2020; Tudi et al., 2022). This study specifically concentrates on the ecological effects of pesticides.

Pesticides can exert their influence over considerable distances from the application site. Those that bind to soil particles may be carried into streams through runoff, and pesticide drift can extend for many miles in the wind. Some pesticides exhibit persistent characteristics, lasting in the environment long after their initial application and posing risks to living organisms over an extended period (Plimmer, 1990). Notably, pesticides such as DDT and chlordane resist easy breakdown and linger in the application site, contributing to soil infertility.

Plants absorb pesticides through their roots and leaves, making them susceptible to the diverse impacts of these chemicals (Alengebawy et al., 2021; Bondareva & Fedorova, 2021). The effects of pesticides on individual plants and animals are varied and multifaceted, underlining the complexity of their ecological repercussions. The longevity of certain pesticides emphasizes the importance of understanding and mitigating their potential adverse effects on the environment.

Conclusion

When deciding whether to use pesticides, it's essential to balance the potential benefits against the associated costs, considering environmental impacts. Despite adhering to a pesticide's label instructions, there is a possibility of causing harm. In such cases, users must carefully weigh the costs and benefits, and if necessary, opt not to use the product.

For those choosing to use pesticides, it is crucial to read and follow the label instructions precisely. While this reduces the risk, it does not guarantee the prevention of accidents. Careful consideration of the environmental hazards section is necessary to assess potential harm to living organisms. If there is a risk, opting for an alternative product is advisable, preferably choosing the least toxic option that effectively addresses the issue. Integrated pest management (IPM) offers a viable alternative, potentially requiring fewer or no pesticides.

Knowing your application area well is key. If specific spots are crucial for unique plants or wildlife, proactive measures should be taken to prevent contamination, especially during critical times of the year. Instead of resorting to pesticides, organic farming stands out as an optimal approach for maintaining a healthy environment. Abstaining from pesticides contributes to soil fertility preservation and serves as a preventive measure for human health.

Table 1. Impact of pesticide on the different insect pest in relation with seasons and crops

Name of the crop	Season	Pest	Pesticide	Chemical compound used
<i>Daucus carota</i> <i>Subsp.sativus</i> (carrot)	July- February	Vegetable weaver/ aphids	linuron	400 g/litre linuron SC
<i>Brassica oleracea</i> <i>var. capitata</i> (cabbage)	October- November	Diamond back moth	FAME	Chlorantran Iliprate 18.5% SC
<i>Phaseolus vulgaris</i> (Beans)	July- February	Pod borer/beetle String bean insect	carbaryl	Carbaryl 43% (1- naphthyl methyl carbamate)
<i>Oryza sativa</i> (Paddy)	November- January	Spotted boll worm	FAME	Chlorantran Iliprate 18.5% SC
<i>Manihot esculenta</i> (Tapioca)	April-may	whitefly	NANMA	Neem oil as big formulation
<i>Allium sepa</i> (Onion)	April-July	Aphid	confidor	Imidacloprid 350 SC
<i>Beta vulgaris</i> (Beetroot)	July- august	Beet leafhopper/ Boll worm	PROFEX	Profenofos 40%+ cypermethrin 4%E.C
<i>Solanum</i> <i>melongena</i> (Brinjal)	June- September	Shoot and root borer/ jassids	carbaryl	Carbaryl 43% (1- naphthyl methyl carbamate)
<i>Abelmoscus</i> <i>esculentus</i> (Ladies finger)	June – August	Shoot and root borers/aphids/ flea/green stinkbugs	Roker	Profenofos and pyrethroid cypermethrin
<i>Solanum</i> <i>lycopersicum</i> (tomato)	March - June	Root- knot nematode	Abamectin	80% avermectin B _{1a} and 20% avermectin B ₁
<i>Raphanus sativus</i> (Radish)	March- august	Aphids	Malathion	Dimethyl dithiophosphoric acid and diethyl fumarate
<i>Trichosanthes</i> <i>cucumerin</i> (snake gourd)	January - march/ September - December	Fruit fly	Dimethoate or neem oil	Dimethoate 400E.C
<i>Lagenaria</i> <i>siceraria</i> (bottle guard)	January - march/ September- December	Squash bug/ whitefly	Neem oil or soap oil	Azadiractin and nimbin
<i>Helianthus annus</i> (sunflower)	January-June	Cutworms/ sunflower beetle	Soap or neem oil	Azadiractin and nimbin 46% nitrogen
<i>Gossypium</i> <i>barbadense</i> (cotton)	April -may/ December - January	Bollworm/ fruit worm	TRACK	Profenofos 40% and cypermethrin 4% E.C
<i>Ipomoea batatas</i> (sweet potato)	June - July	Potato tubeworms/ flea beetles	Pyrethrin	permethrin
<i>Musa acuminata</i> (raw banana)	February – April/	Banana weevil	chlorpyrifos	Chlorinated phosphates

	November-December			
<i>Momordica charantia</i> (bitter gourd)	January - march	Cucurbit fruit fly	Diptrex (trichlorfon)	500 g/l trichlorfon
<i>Luffa</i> (ridge gourd)	January - April	Leaf miner/fruit fly	malathion	Dimethyl dithiophosphoric acid and diethyl fumarate
<i>Triticum aestivum</i> (wheat)	October – December/ February - may	Aphid	Organo chlorides	Chlorines include DDT
<i>Zea mays</i> (maize)	June - July	Stem borer/ shoot fly	glyphosate	Isopropylamine salt
<i>Solanum tuberosum</i> (potato)	October – November	Black and yellow striped potato bud	coragen	Chlorano tranilprole S.C
<i>piper betle</i> (betel)	January - february	Red spider mite/ Aphids/shoot bugs/white fly	fizimite	permethri
<i>Moringa olifera</i> (drumstick)	May - june	Bud worm/ hairy caterpillar	dimethoate	Dimethoate 400E.C
<i>Mangifera indica</i> (mango)	February-march	Mango hopper/fruit fly/stem borer	Dimethoate	Dimethoate 400E.C
<i>Cocus nucifera</i> (coconut)	April	Red palm weevil	Imidacloprid	chloronicotiny
<i>Coccinia grandis</i> (ivy gourd)	February - march	Aphids/ whitefly	Garlon 4	Triclopyr D
<i>Dioscorea alata</i> (elephant foot yam)	April - may	mealybugs	Cow dung slurry with neem cake oil	Azadirachtin and animal waste
<i>Pisum sativum</i> (pea)	February - march/ October – november	Aphids / pea weevil	neonicotinoid	Acetamaprid and thiamethoxam
<i>Spinacia oleracea</i> (spinach)	March- may	Cutworms/ caterpillars	Dimethoate	Dimethoate 400E.C

Acknowledgement

The authors' thankful to the Sri Parasakthi College for Women, Tenkasi, Tamil Nadu, India for providing necessary facilities for carried out of this work.

References

- Abdollahdokht, D., Gao, Y., Faramarz, S., Poustforoosh, A., Abbasi, M., Asadikaram, G., & Nematollahi, M. H. (2022). Conventional agrochemicals towards nano-biopesticides: an overview on recent advances. *Chemical and Biological Technologies in Agriculture*, 9(1), 13. <https://doi.org/10.1186/s40538-021-00281-0>
- Alengebawy, A., Abdelkhalek, S. T., Qureshi, S. R., & Wang, M.-Q. (2021). Heavy Metals and Pesticides Toxicity in Agricultural Soil and Plants: Ecological Risks and Human Health Implications. *Toxics*, 9(3). <https://doi.org/10.3390/toxics9030042>
- Ayilara, M. S., Adeleke, B. S., Akinola, S. A., Fayose, C. A., Adeyemi, U. T., Gbadegesin, L. A., Omole, R. K., Johnson, R. M., Uthman, Q.

- O., & Babalola, O. O. (2023). Biopesticides as a promising alternative to synthetic pesticides: A case for microbial pesticides, phytopesticides, and nanobiopesticides. *Frontiers in Microbiology*, 14, 1040901. <https://doi.org/10.3389/fmicb.2023.1040901>
- Bondareva, L., & Fedorova, N. (2021). Pesticides: Behavior in Agricultural Soil and Plants. *Molecules (Basel, Switzerland)*, 26(17). <https://doi.org/10.3390/molecules26175370>
- Cooper, J., & Dobson, H. (2007). The benefits of pesticides to mankind and the environment. *Crop Protection*, 26(9), 1337–1348. <https://doi.org/https://doi.org/10.1016/j.cropro.2007.03.022>
- Damalas, C. A., & Eleftherohorinos, I. G. (2011). Pesticide exposure, safety issues, and risk assessment indicators. *International Journal of Environmental Research and Public Health*, 8(5), 1402–1419. <https://doi.org/10.3390/ijerph8051402>
- Dhanaraju, M., Chenniappan, P., Ramalingam, K., Pazhanivelan, S., & Kaliaperumal, R. (2022). Smart Farming: Internet of Things (IoT)-Based Sustainable Agriculture. *Agriculture*, 12(10). <https://doi.org/10.3390/agriculture12101745>
- Fleeger, J. W. (2020). How Do Indirect Effects of Contaminants Inform Ecotoxicology? A Review. *Processes*, 8(12). <https://doi.org/10.3390/pr8121659>
- Hassaan, M. A., & El Nemr, A. (2020). Pesticides pollution: Classifications, human health impact, extraction and treatment techniques. *The Egyptian Journal of Aquatic Research*, 46(3), 207–220. <https://doi.org/https://doi.org/10.1016/j.ejar.2020.08.007>
- Nicolopoulou-Stamati, P., Maipas, S., Kotampasi, C., Stamatis, P., & Hens, L. (2016). Chemical Pesticides and Human Health: The Urgent Need for a New Concept in Agriculture. *Frontiers in Public Health*, 4, 148. <https://doi.org/10.3389/fpubh.2016.00148>
- Park, D. W., Yang, Y. S., Lee, Y.-U., Han, S. J., Kim, H. J., Kim, S.-H., Kim, J. P., Cho, S. J., Lee, D., Song, N., Han, Y., Hee Kim, H., Cho, B.-S., Chung, J. K., & Kim, A. G. (2021). Pesticide Residues and Risk Assessment from Monitoring Programs in the Largest Production Area of Leafy Vegetables in South Korea: A 15-Year Study. *Foods (Basel, Switzerland)*, 10(2). <https://doi.org/10.3390/foods10020425>
- Pathak, V. M., Verma, V. K., Rawat, B. S., Kaur, B., Babu, N., Sharma, A., Dewali, S., Yadav, M., Kumari, R., Singh, S., Mohapatra, A., Pandey, V., Rana, N., & Cunill, J. M. (2022). Current status of pesticide effects on environment, human health and it's eco-friendly management as bioremediation: A comprehensive review. *Frontiers in Microbiology*, 13(August), 1–29. <https://doi.org/10.3389/fmicb.2022.962619>
- Plimmer, J. R. (1990). Pesticide loss to the atmosphere. *American Journal of Industrial Medicine*, 18(4), 461–466. <https://doi.org/10.1002/ajim.4700180418>
- Ratnadass, A., Fernandes, P., Avelino, J., & Habib, R. (2012). Plant species diversity for sustainable management of crop pests and diseases in agroecosystems: a review. *Agronomy for Sustainable Development*, 32(1), 273–303. <https://doi.org/10.1007/s13593-011-0022-4>
- Skendžić, S., Zovko, M., Živković, I. P., Lešić, V., & Lemić, D. (2021). The Impact of Climate Change on Agricultural Insect Pests. *Insects*, 12(5). <https://doi.org/10.3390/insects12050440>
- Tudi, M., Daniel Ruan, H., Wang, L., Lyu, J., Sadler, R., Connell, D., Chu, C., & Phung, D. T. (2021). Agriculture Development, Pesticide Application and Its Impact on the Environment. *International Journal of Environmental Research and Public Health*, 18(3). <https://doi.org/10.3390/ijerph18031112>
- Tudi, M., Li, H., Li, H., Wang, L., Lyu, J., Yang, L., Tong, S., Yu, Q. J., Ruan, H. D., Atabila, A., Phung, D. T., Sadler, R., & Connell, D. (2022). Exposure Routes and Health Risks Associated with Pesticide Application. *Toxics*, 10(6). <https://doi.org/10.3390/toxics10060335>